



Orchid Disease Control

Part 1 - How Orchids Respond to Disease Pathogens

by Sue Bottom, sbottom15@hotmail.com

The best way to prevent orchid disease is to grow strong healthy plants that are able to use their natural defense mechanisms to protect themselves. Orchids don't have an immunological system like we do. They have a phloem that transports the sugars produced in the leaves by photosynthesis through the rest of the plant and a xylem that transports water and soluble mineral nutrients absorbed from the roots upwards throughout the plant, but they don't have a circulatory system carrying leukocytes, lymphocytes, antibodies and all the rest of our incredible infection fighters. So how does an orchid protect itself from disease?



Black Rot on Cattleya - Supplying your plants with sufficient calcium will strengthen them so they can more easily withstand attack by the water molds that cause Black Rot though those suffering through hot humid summers may also have to rely on heavy duty fungicides to protect their plants.

The Natural Disease Response in Orchids – Wall Off the Invading Pathogens

Barrier to Entry. Orchids have an epidermis with an outer layer called a cuticle that consists of a wax-like substance that provides protection against water loss as well as a barrier to entry, acting as the first line of defense against pathogens. Some fungi can pierce this barrier by secreting an enzyme that degrades the cuticle. Bacteria and some fungi circumvent the barrier by entering tissues through open stomata or wound sites.

Hypersensitive Response. When an attack by a pathogen begins, plants respond with a rapid series of events at the site of the infection that culminates in the hypersensitive response. The hypersensitive response is a localized death of cells at the site of infection as a result of the plant cells recognizing the invading pathogen and committing suicide to thwart the pathogen. The dead cells wall off the pathogen and this necrotic barrier prevents the transmission of the pathogen through the plant. These dead cells may also become hard and unattractive to other pathogens by producing toxic compounds.



St. Augustine Orchid Society

www.staugorchidsociety.org

Orchid Disease Control


Part 1 - How Orchids Respond to Disease Pathogens

by Sue Bottom, sbottom15@hotmail.com

Systemic Acquired Resistance. The initial Hypersensitive Response is commonly followed by a slower response that leads to Systemic Acquired Resistance (SAR). Systemic Acquired Resistance occurs when a chemical travels from the infection site to nearby tissues via the phloem and primes the rest of the plant to protect itself against further challenges from the invading pathogen. The plant's response to the infection in one leaf causes the plant to become more resistant to a secondary attack in a different part of the plant.

Substances that May Induce a Plant's Response to Disease – Artificial Stimulation of the Plant's Response Mechanism

Aspirin. Salicylic acid (a chemical related to aspirin) is one of several chemicals that can transmit the signal of the defense response throughout the plant. Studies have shown that applying salicylic acid to plants can induce resistance to pathogens, environmental stresses and some insects. Well known Ft. Lauderdale orchid grower Dot Henley wrote about her experience applying aspirin to her orchids in the October 2001 edition of the AOS Bulletin. She reported that her plants had more flowers, bigger growths and fewer fungal problems after weekly spraying with a fertilizer solution to which aspirin was added (3/4 of a 325 mg aspirin in a gallon of water). She later added two caveats, do not exceed the recommended amount and do not skip adding aspirin to your fertilizer regimen for more than a couple of weeks because the orchids seem to become addicted to aspirin and fungal and bacterial problems may occur without their weekly fix.



Occasional preventative treatments can help. I spray out of bloom phalaenopsis and paphiopedilums twice a year with copper compounds, after repotting in the summer and before the cool season in the fall to help prevent bacterial infections from *Erwinia* (now *Pectobacterium*).

Harpin Proteins. A naturally occurring protein isolated from bacterial plant pathogens has been shown to initiate the protective response in plants when applied to leaves. Plants can detect the presence of these Harpin proteins and their early warning system is activated. The plant thinks it is under attack and signals other parts of the plant to activate defensive and growth responses. Rx Green Solutions markets Axiom, the Harpin Alpha Beta Protein, and recommends it be applied every 2 to 3 weeks at a rate of a 2 gram packet per gallon as a foliar application via a light mist. The potential effectiveness of Harpin proteins for orchids is currently under evaluation in the scientific community.



St. Augustine Orchid Society

www.staugorchidsociety.org

Orchid Disease Control

Part 1 - How Orchids Respond to Disease Pathogens

by Sue Bottom, sbottom15@hotmail.com

Biological Agents. Biological control agents are introduced or resident living organisms that suppress the activities and populations of plant pathogens. They are most effective if applied prior to infection. Biological control agents may work by killing the pathogen, producing a compound that is toxic or interferes with pathogen growth, triggering a defensive response in the host plant or outcompeting the pathogen. Promix HP has a biofungicide incorporated into their soilless mix, a naturally occurring strain of bacteria (*Bacillus subtilis*, *Bacillus pumilus*) reported to enhance plant growth and suppress harmful organisms such as *Fusarium*, *Pythium* and *Rhizoctonia*. There are a variety of other products containing *Streptomyces lydicus*, *Bacillus subtilis*, *Trichoderma* species and *Gliocladium virens* that claim to help prevent some soil borne diseases and aerial diseases if applied prior to the disease occurring in the growing area.

Wound Management and Protective Sprays – Kill Pathogens in the Environment Before They Infect Your Plant. There are a raft of chemicals available to protect your plants from bacterial and fungal infections although they are most effective as protectants rather than curatives. There are many precautionary treatments you can incorporate into your growing regimen. Regularly disinfection of your growing area by spraying with sterilants like bleach or quaternary ammonium compounds is a good start. Many orchid growers dust open wounds on their plants with cinnamon or a fungicide to seal wounds and prevent infection. After repotting, I pour a protective drench of Banrot through the fresh potting mix to ward off infection, plus some seaweed extract to encourage root growth. Some spray specific types of plants for a specific potential problem, like monthly sprays with thiophanate methyl to control *Guignardia* in vandas and a post repotting and fall spray of copper compounds to help prevent *Erwinia* (now called *Pectobacterium*) in phals and paphs.



Good air movement is one of the most effective ways to prevent a disease from infecting your plant, like this *Cercospora* damage on a cattleya. Fresh air constantly bathing the leaves together with good spacing between plants will help prevent a disease from gaining a stronghold in your growing area.

Good Cultural Practices – Your Best Defense Against Orchid Disease is a Vibrantly Growing Plant that Can Resist Infection. Your best defense against orchid disease is to optimize your culture so your orchids are healthy and happy and able to resist diseases naturally. This means you must provide your plants with enough of the right kinds of light, water, nutrients, warmth and air movement and make sure your potting media is fresh. The importance of fresh air bathing your plants cannot be overemphasized. Fresh air will only work its wonders if there is sufficient spacing between plants. Sanitation is vitally important to eliminate the source of pathogens in your growing area. Maintain a clean growing area free of weeds that often



St. Augustine Orchid Society

www.staugorchidsociety.org

Orchid Disease Control

Part 1 - How Orchids Respond to Disease Pathogens

by Sue Bottom, sbottom15@hotmail.com

harbor disease. Remove spent flowers and fallen leaves from the growing area to a sealed container to remove the source of infections. Disinfect benches and under benches regularly to kill any lingering pathogens. Sterilize cutting tools, pots, wire products, etc. before reuse so problems aren't transferred from one plant to another. Always use new or sterile potting medium. Don't allow water to drip from one plant to another and never dunk plants in a common bucket. Make sure your new orchids are healthy, happy and disease free before introducing them to your growing area. When you water, look at each plant for any signs of stress, particularly as seasonal changes in temperature, light and moisture occur. Consider developing a schedule for preventative applications of fungicides or bactericides suitable to your climate and growing conditions.



Don't be this grower! Never allow spent flowers or vegetation to linger in your growing area. Spores and other inoculum for future infections are growing on these dead plant parts. Promptly remove them from the growing area to a sealed container for disposal, no composting!

Many disease problems can be prevented with attention to sanitation and good cultural practices. Occasionally you may find you have a genetic weakling that is prone to problems. I tend to discard these unless the flower is extraordinary. All growers will find disease in their collections from time to time and have to resort to the use of fungicides



St. Augustine Orchid Society

www.staugorchidsociety.org

Orchid Disease Control

Part 1 - How Orchids Respond to Disease Pathogens

by Sue Bottom, sbottom15@hotmail.com

and bactericides to control a problem. If you have a recurring disease problem, there may be some aspect of your cultural practices that should be changed or perhaps you are trying to grow a plant not well suited to your growing conditions. Your plants will be happiest if you focus on improving your culture rather than reflexively spraying chemicals on them whenever you notice a new leaf spot.

Citations and Additional Reading:

Chuang, H. W., Chang, P. Y., & Syu, Y. Y. Harpin Protein, an Elicitor of Disease Resistance, Acts as a Growth Promoter in Phalaenopsis Orchids. *Journal of Plant Growth Regulation*, 1-10.

Freeman, B.C. and G.A. Beattie. 2008. An Overview of Plant Defenses against Pathogens and Herbivores. *The Plant Health Instructor*. Accessed online
<http://www.apsnet.org/edcenter/intropp/topics/Pages/OverviewOfPlantDiseases.aspx>

Freeman, S., Quillin, K. and Allison, L., Chapter 37, Plant Defense Systems, pp. 709-725 Biological Science, 4th Edition, Prentice Hall. http://www.pearsoncustom.com/ucsc/images/0130819239_ch37.pdf, Accessed May 26, 2014

Guest, D., & Brown, J., 1997. Plant defences against pathogens. *Plant Pathogens and Plant Diseases*, 263-286. [http://www.apsnet.org/Publications/Brown_Ogle/17_Defence_mechanisms_\(DIG&JFB\).pdf](http://www.apsnet.org/Publications/Brown_Ogle/17_Defence_mechanisms_(DIG&JFB).pdf) Accessed May 26, 2014

Harmon, P.F., Bledsoe, S.D. and McGovern, R.J., Professional Disease Management Guide for Ornamental Plants, University of Florida, PP-202, August 2004, Revised 2012

Henley, D., Aspirin and Growing Orchids, *Orchids*, 70(10). 983.

Pal, K. K. and Gardener, B.M., 2006. Biological Control of Plant Pathogens. *The Plant Health Instructor* 2, pp. 1117-1142

Raskin, I., 1992. Salicylate, A New Plant Hormone. *Plant Physiology*, July 1992 pp. 799-803.